

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

Warnings

- Do not use**
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
 - right before or after heart surgery
- Ask a doctor before use if you have**
- never had migraines diagnosed by a health professional
 - a headache that is different from your usual migraines
 - the worst headache of your life
 - fever and stiff neck
 - headaches beginning after or caused by head injury, exertion, coughing or bending
 - experienced your first headache after the age of 50
 - daily headaches
 - a migraine so severe as to require bed rest
 - problems or serious side effects from taking pain relievers or fever reducers
 - stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
 - ulcers
 - bleeding problems
 - high blood pressure
 - heart or kidney disease
 - asthma
 - taken a diuretic
 - reached age 60 or older
- Ask a doctor or pharmacist before use if you are**
- taking any other drug containing an NSAID (prescription or nonprescription)
 - taking a blood thinning (anticoagulant) or steroid drug
 - under a doctor's care for any serious condition
 - taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 - taking any other drug
- When using this product**
- take with food or milk if stomach upset occurs
 - the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
- Stop use and ask a doctor if**
- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 - migraine headache pain is not relieved or gets worse after first dose
 - stomach pain or upset gets worse or lasts
 - any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Other information

- treats migraine
- Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin reddening
 - rash
 - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.
- This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- do not take more than directed**
- the smallest effective dose should be used**

adults	<ul style="list-style-type: none">• take 2 capsules with a glass of water• if symptoms persist or worsen, ask your doctor• do not take more than 2 capsules in 24 hours, unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">• ask a doctor

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

D&C yellow no. 10, FD&C green no. 3, FD&C red no. 40, gelatin, light mineral oil, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol